

1  
2  
3  
4  
5  
6 **COMMODITY SPECIFIC FOOD SAFETY GUIDELINES FOR THE**  
7 **PRODUCTION AND HARVEST OF LETTUCE AND LEAFY GREENS**  
8 **VERSION 2 - ARIZONA**  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

**AUGUST 26, 2008**

38 Authors Note:

39 This document reflects Commodity Specific Food Safety Guidelines for the  
40 Production and Harvest of Leafy Greens for Arizona.. It is based on the Commodity  
41 Specific Food Safety Guidelines for the Production and Harvest of Leafy Greens  
42 accepted for use by the California Leafy Greens Handler Marketing Agreement and  
43 contains minor, non-substantive modifications recommended by the Arizona Leafy  
44 Greens Marketing Committee. Arizona law supersedes any requirements in this  
document that may be in conflict.

45 **TABLE OF CONTENTS**

46

47 Glossary ..... 3

48 Acronyms and Abbreviations ..... 7

49 List of Appendices ..... 8

50 Introduction..... 9

51 Scope..... 10

52 1. Purpose..... 12

53 2. Issue: General Requirements ..... 12

54 3. Issue: Environmental Assessments ..... 12

55 4. Issue: Water ..... 13

56 5. Issue: Soil Amendments ..... 23

57 6. Issue: Nonsynthetic Crop Treatments..... 32

58 7. Issue: Harvest Equipment ..... 36

59 8. Issue: Harvest Personnel - Direct Contact with Soil during Harvest..... 38

60 9. Issue: Field Personnel - Transfer of Human Pathogens by Field Workers..... 38

61 10. Issue: Equipment Facilitated Cross Contamination..... 39

62 11. Issue: Flooding..... 40

63 12. Issue: Water Usage to Prevent Product Dehydration..... **Error! Bookmark not**

64 **defined.**

65 13. Issue: Production Locations - Climatic Conditions and Environment..... 44

66 14. Issue: Production Locations - Encroachment by Animals and Urban Settings .... 44

67 15. Detailed Background Guidance Information ..... 53

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

<b>Aerosolized</b>	The dispersion or discharge of a substance under pressure that generates a suspension of fine particles in air or other gas.
<b>animal by-product</b>	Most parts of an animal that do not include muscle meat including organ meat, nervous tissue, cartilage, bone, blood and excrement.
<b>animals of significant risk</b>	Animals that have been determined by the Centers for Disease Control to have a higher risk of carrying E. coli O157:H7. These animals are cattle, sheep, goats, pigs (domestic and wild), and deer.
<b>adenosine tri-phosphate (ATP)</b>	A high energy phosphate molecule required to provide energy for cellular function.
<b>ATP test methods</b>	Exploits knowledge of the concentration of ATP as related to viable biomass or metabolic activity; provides an estimate of cleanliness.
<b>Biofertilizers</b>	Fertilizer materials/products that contain microorganisms such as bacteria, fungi, and cyanobacteria that shall promote soil biological activities..
<b>Biosolids</b>	Solid, semisolid, or liquid residues generated during primary, secondary, or advanced treatment of domestic sanitary sewage through one or more controlled processes.
<b>colony forming units (CFU)</b>	Viable micro-organisms (bacteria, yeasts & mold) either consisting of single cells or groups of cells, capable of growth under the prescribed conditions (medium, atmosphere, time and temperature) to develop into visible colonies (colony forming units) which are counted.
<b>Concentrated Animal Feeding Operation (CAFO)</b>	A lot or facility where animals have been, are or will be stabled or confined and fed or maintained for a total of 45 days or more in any 12 month period and crops, vegetation forage growth, or post-harvest residues are not sustained in the normal growing season over any portion of the lot or facility. In addition, there must be more than 1,000 'animal units' (as defined in 40 CFR 122.23) confined at the facility; or more than 300 animal units confined at the facility if either one of the following conditions are met: pollutants are discharged into navigable waters through a man-made ditch, flushing system or other similar man-made device; or pollutants are discharged directly into waters of the United States which originate outside of and pass over, across, or through the facility or otherwise come into direct contact with the animals confined in the

	operation.
<b>coliforms</b>	Gram-negative, non-sporeforming, rod-shaped bacteria that ferment lactose to gas. They are frequently used as indicators of process control, but exist broadly in nature.
<b>cross contamination</b>	The transfer of microorganisms, such as bacteria and viruses, from one place to another.
<b>E. coli</b>	<i>Escherichia coli</i> is a common bacteria that lives in the lower intestines of animals (including humans) and is generally not harmful. It is frequently used as an indicator of fecal contamination, but can be found in nature from non-fecal sources.
<b>fecal coliforms</b>	Coliform bacteria that grow at elevated temperatures and may or may not be of fecal origin. Useful to monitor effectiveness of composting processes. Also called “thermotolerant coliforms.”
<b>Flooding</b>	The flowing or overflowing of a field with water outside a producer’s control that is reasonably likely to contain microorganisms of significant public health concern and is reasonably likely to cause adulteration of edible portions of fresh produce in that field.
<b>food contact surface</b>	A surface of equipment or a utensil with which food normally comes into contact, or from which food may drain, drip or splash into a food or onto a surface normally in contact with food.
<b>food safety assessment</b>	A standardized procedure that predicts the likelihood of harm resulting from exposure to chemical, microbial and physical agents in the diet.
<b>food safety professional</b>	Person entrusted with management level responsibility for conducting food safety assessments before food reaches consumers; requires formal training in scientific principles and a solid understanding of the principles of food safety as applied to agricultural production.
<b>geometric mean</b>	Mathematical def.: the n-th root of the product of n numbers, or: Geometric Mean = n-th root of $(X_1)(X_2)...(X_n)$ , where $X_1, X_2,$ etc. represent the individual data points, and n is the total number of data points used in the calculation. Practical def.: the average of the logarithmic values of a data set, converted back to a base

	10 number.
<b>Hydroponic</b>	The growing of plants in nutrient solutions with or without an inert medium (as soil) to provide mechanical support.
<b>Indicator microorganisms</b>	An organism that when present suggests the possibility of contamination or under processing.
<b>leafy greens</b>	Iceberg lettuce, romaine lettuce, green leaf lettuce, red leaf lettuce, butter lettuce, baby leaf lettuce (i.e., immature lettuce or leafy greens), escarole, endive, spring mix, spinach, cabbage (green, red and savoy), kale, arugula and chard.
<b>Monthly</b>	Because irrigation schedules and delivery of water is not always in a growers control “monthly” for purposes of water sampling means within 35 days of the previous sample.
<b>most probable number (MPN)</b>	Estimated values that are statistical in nature; a method for enumeration of microbes in a sample, particularly when present in small numbers.
<b>nonsynthetic crop treatments</b>	Any crop input that contains animal manure, an animal product, and/or an animal by-product that is reasonably likely to contain human pathogens.
<b>Ready to eat (RTE) food</b> <i>(excerpted from USFDA 2005 Model Food Code)</i>	(1) "Ready-to-eat food" means FOOD that: (a) Is in a form that is edible without additional preparation to achieve FOOD safety, as specified under one of the following: 3-401.11(A) or (B), § 3-401.12, or § 3-402.11, or as specified in 3-401.11(C); or (d) May receive additional preparation for palatability or aesthetic, epicurean, gastronomic, or culinary purposes. (2) "Ready-to-eat food" includes: (b) Raw fruits and vegetables that are washed as specified under § 3-302.15; (c) Fruits and vegetables that are cooked for hot holding, as specified under § 3-401.13; (e) Plant FOOD for which further washing, cooking, or other processing is not required for FOOD safety, and from which rinds, peels, husks, or shells, if naturally present are removed;
<b>synthetic crop treatments (chemical fertilizers)</b>	Any crop inputs that may be refined, and/or chemically synthesized and/or transformed through a chemical process (e.g. gypsum, lime, sulfur, potash, ammonium sulfate etc.).
<b>oxidation reduction potential (ORP)</b>	An intrinsic property that indicates the tendency of a chemical species to acquire electrons and so be reduced; the more positive

	the ORP, the greater the species' affinity for electrons.
<b>parts per million (ppm)</b>	Usually describes the concentration of something in water or soil; one particle of a given substance for every 999,999 other particles.
<b>Pathogen</b>	A disease causing agent such as a virus, parasite, or bacteria.
<b>pooled water</b>	An accumulation of standing water; not free-flowing.
<b>process authority</b>	A regulatory body, person, or organization that has specific responsibility and knowledge regarding a particular process or method; these authorities publish standards, metrics, or guidance for these processes and/or methods.
<b>risk mitigation</b>	actions to reduce the severity/impact of a risk
<b>soil amendment</b>	Elements added to the soil, such as compost, peat moss, or fertilizer, to improve its capacity to support plant life.
<b>ultraviolet index (UV index)</b>	A measure of the solar ultraviolet intensity at the Earth's surface; indicates the day's exposure to ultraviolet rays. The UV index is measured around noon for a one-hour period and rated on a scale of 0-15.
<b>Validated process</b>	A process that has been demonstrated to be effective through a statistically-based study, literature, or regulatory guidance.
<b>water distribution system</b>	Distribution systems -- consisting of pipes, pumps, valves, storage tanks, reservoirs, meters, fittings, and other hydraulic appurtenances -- canals, ditches and rivers -- to carry water from its primary source to a lettuce and leafy green crop.

92

93

94	<b>ACRONYMS AND ABBREVIATIONS</b>
95	
96	AFOs: Animal feeding operations
97	AOAC: the Association of Official Agricultural Chemists
98	BAM: Bacteriological Analytical Manual
99	CAFOs: Concentrated animal feeding operations
100	CSG2: <i>Commodity Specific Guidance for Leafy Greens and Lettuce, 2<sup>nd</sup> Edition</i>
101	CFU: colony forming units
102	cGMP: current good manufacturing practices
103	COA: Certificate of Analysis
104	DL: Detection Limit
105	FDA: Food and Drug Administration
106	GAPS: good agricultural practices
107	GLPs: good laboratory practices
108	HACCP: hazard analysis critical control point
109	MPN: most probable number
110	NGO: nongovernmental organization
111	NRCS: Natural Resources Conservation Service
112	ORP: Oxidation reduction potential
113	PPM: parts per million
114	RTE: ready-to-eat
115	SSOPs: Sanitation Standard Operating Procedures
116	USEPA: United States Environmental Protection Agency
117	UV: ultraviolet
118	WHO: World Health Organization
119	
120	
121	
122	
123	
124	
125	
126	
127	
128	
129	

130	<b>LIST OF APPENDICES</b>	
131	<a href="#">Appendix A</a> : Sanitary Survey	<a href="#">Appendix B</a> : Technical Basis Document
132	<a href="#">Appendix C</a> : Crop Sampling Protocol	<a href="#">Appendix Z</a> : Resource Agency Contacts
133	<a href="#">Appendix D</a> : Kinetics of Microbial Inactivation for Alternative Food Processing Technologies	
134		

135 **INTRODUCTION**

136

137 In 1998, the U.S. Food and Drug Administration (FDA) issued its “Guide to Minimize  
138 Microbial Food Safety Hazards for Fresh Fruits and Vegetables.” The practices outlined in  
139 this and other industry documents are collectively known as Good Agricultural Practices or  
140 GAPs. GAPs provide general food safety guidance on critical production steps where food  
141 safety might be compromised during the growing, harvesting, transportation, cooling,  
142 packing and storage of fresh produce. More specifically, GAP guidance alerts fruit and  
143 vegetable producers, shippers, packers and processors to the potential microbiological  
144 hazards associated with various aspects of the production chain including: land history,  
145 adjacent land use, water quality, worker hygiene, pesticide and fertilizer use, equipment  
146 sanitation and product transportation. The vast majority of the lettuce/leafy greens industry  
147 has adopted GAPs as part of normal production operations. Indeed the majority of  
148 lettuce/leafy greens producers undergo either internal or external third-party GAP audits on a  
149 regular basis to monitor and verify adherence to their GAPs programs. These audit results are  
150 often shared with customers as verification of the producer’s commitment to food safety and  
151 GAPs.

152

153 While the produce industry has an admirable record of providing the general public with safe,  
154 nutritious fruits and vegetables, it remains committed to continuous improvement with regard  
155 to food safety. In 2004, the FDA published a food safety action plan that specifically  
156 requested produce industry leadership in developing the next generation of food safety  
157 guidance for fruit and vegetable production. These new commodity-specific guidelines focus  
158 on providing guidance that enhances the safe growing, processing, distribution and handling  
159 of commodities from the field to the end user. The 1<sup>st</sup> Edition of these new voluntary  
160 guidelines were published by the industry in April 2006.

161 In response to continued concerns regarding the microbial safety of fresh produce, this  
162 edition of the guidelines (which focuses solely on production and harvest practices) was  
163 prepared to provide more specific and quantitative measures of identified best practices. A  
164 key focus of this revision was to identify, where possible and practical, metrics and measures  
165 that could be used to assist the industry with compliance with the guidelines. In preparing  
166 this document, metrics were researched for three primary areas: water quality, soil  
167 amendments, and environmental assessments/conditions. A three-tier approach was used to  
168 identify these metrics in as rigorous a manner as possible:

- 169 1. A comprehensive literature review was conducted to determine if there was a  
170 scientifically valid basis for establishing a metric for the identified risk factor or best  
171 practice.
- 172 2. If the literature research did not identify scientific studies that could support an  
173 appropriate metric, standards or metrics from authoritative or regulatory bodies were  
174 used to establish a metric.
- 175 3. If neither scientific studies nor authoritative bodies had allowed for suitable metrics,  
176 consensus among industry representatives and/or other stakeholders was sought to  
177 establish metrics.

178 In the last 10 years, the focus of food safety efforts has been on the farm, initial cooling and  
179 distribution points, and value-added processing operations. Fruit and vegetable processing

180 operations have developed sophisticated food safety programs largely centered on current  
181 Good Manufacturing Practices (cGMPs) and the principles of Hazard Analysis Critical  
182 Control Point (HACCP) programs. As we develop a greater understanding of food safety  
183 issues relative to the full spectrum of supply and distribution channels for fruits and  
184 vegetables, it has become clear that the next generation of food safety guidance needs to  
185 encompass the entire supply chain.

186 In addition to this document, several supplemental documents have been prepared to explain  
187 the rationale for the metrics and assist the producer with activities in the field. These  
188 documents include a “Technical Basis Document” that describes in detail and with  
189 appropriate citations the bases for the changes made in this edition of this document, a  
190 Sanitary Survey document that describes the processes for assessing the integrity and  
191 remediation of water systems, and an example product testing plan. All of these items can be  
192 found as Appendices to this document.

### 193 **SCOPE**

194 The scope of this document pertains only to fresh and fresh-cut lettuce and leafy greens  
195 products. It does not include products commingled with non-produce ingredients (e.g. salad  
196 kits which may contain meat, cheese, and/or dressings). Examples of “lettuce/leafy greens”  
197 include iceberg lettuce, romaine lettuce, green leaf lettuce, red leaf lettuce, butter lettuce,  
198 baby leaf lettuce (i.e., immature lettuce or leafy greens), escarole, endive, spring mix, cabbage  
199 (green, red and savoy), kale, arugula, chard and spinach. These crops are typically considered  
200 lettuce and leafy greens by FDA but may not be similarly defined by other state or federal  
201 regulatory bodies. This document is also limited to offering food safety guidance for crops  
202 grown under outdoor field growing practices and may not address food safety issues related  
203 to hydroponic and/or soil-less media production techniques for lettuce/leafy greens.

204 Lettuce/leafy greens may be harvested mechanically or by hand and are almost always  
205 consumed uncooked or raw. Because lettuce/leafy greens may be hand-harvested and hand-  
206 sorted for quality, there are numerous “touch points” early in the supply chain and a similar  
207 number of “touch points” later in the supply chain as the products are used in foodservice or  
208 retail operations. Each of these “touch points” represents a potential opportunity for cross-  
209 contamination. For purposes of this document, a “touch point” is any occasion when the  
210 food is handled by a worker or contacts an equipment food contact surface.

211  
212 Lettuce/leafy greens present multiple opportunities to employ food safety risk management  
213 practices to enhance the safety of lettuce/leafy greens. In the production and harvest of  
214 lettuce and leafy greens as raw agricultural commodities, GAPs are commonly employed in  
215 order to produce the safest products possible. In a processing operation, the basic principles  
216 of cGMPs, HACCP, sanitation and documented operating procedures are commonly  
217 employed in order to produce the safest products possible. Lettuce/leafy greens are highly  
218 perishable and it is strongly recommended that they be distributed, stored and displayed  
219 under refrigeration.

220  
221 Safe production, packing, processing, distribution and handling of lettuce/leafy greens  
222 depend upon a myriad of factors and the diligent efforts and food safety commitment of  
223 many parties throughout the distribution chain. No single resource document can anticipate  
224 every food safety issue or provide answers to all food safety questions. These guidelines

225 focus on minimizing only the microbial food safety hazards by providing suggested actions  
226 to reduce, control or eliminate microbial contamination of lettuce/leafy greens in the field to  
227 fork distribution supply chain.

228 All companies involved in the lettuce/leafy greens farm to table supply chain shall implement  
229 the recommendations contained within these guidelines to provide for the safe production and  
230 handling of lettuce/leafy greens products from field to fork. Every effort to provide food  
231 safety education to supply chain partners should also be made. Together with the  
232 commitment of each party along the supply chain to review and implement these guidelines,  
233 the fresh produce industry is doing its part to provide a consistent, safe supply of produce to  
234 the market.

235  
236 These guidelines are intended only to convey the best practices associated with the industry.  
237 The Produce Marketing Association, the United Fresh Produce Association, Western  
238 Growers, and all other contributors and reviewers make no claims or warranties about any  
239 specific actions contained herein. It is the responsibility of any purveyor of food to maintain  
240 strict compliance with all local, state and federal laws, rules and regulations. These  
241 guidelines are designed to facilitate inquiries and developing information that must be  
242 independently evaluated by all parties with regard to compliance with legal and regulatory  
243 requirements. The providers of this document do not certify compliance with these guidelines  
244 and do not endorse companies or products based upon their use of these guidelines.

245 Differences between products, production processes, distribution and consumption, and the  
246 ever-changing state of knowledge regarding food safety make it impossible for any single  
247 document to be comprehensive and absolutely authoritative. Users of these guidelines should  
248 be aware that scientific and regulatory authorities are periodically revising information  
249 regarding best practices in food handling, as well as information regarding potential food  
250 safety management issues. Users of this document must bear in mind that as knowledge  
251 regarding food safety changes, measures to address those changes will also change as will the  
252 emphasis on particular issues by regulators and the regulations themselves. Neither this  
253 document nor the measures food producers and distributors should take to address food  
254 safety are set in stone.

255 Due to the close association between production blocks and environmentally sensitive areas  
256 in many locations, it is important to consult environmental regulators when any mitigation  
257 strategies that may impact these areas are employed. Producers should implement strategies  
258 that not only protect food safety but also support conservation practices, water quality, and  
259 habitat protection. All parties involved with implementing the practices outlined in this  
260 document should be aware that these metrics are not, in any way, meant to encourage  
261 producers to violate environmental regulations.

262  
263 Users are strongly urged to maintain regular contact with and utilize information available  
264 from their trade associations, the U.S. Food and Drug Administration, the U.S. Department of  
265 Agriculture, the U.S. Environmental Protection Agency, the Centers for Disease Control and  
266 Prevention, and state agricultural, environmental, academic, and public health authorities.

267 The Sanitary Survey and Technical Basis Documents prepared as Appendices to these  
268 guidelines are considered to be additional resources. They are intended to provide  
269 clarification, assist with interpretation and provide additional guidance as users develop food  
270 safety programs based on these Guidelines. They are not intended for measurement or  
271 verification purposes.

272 **Lettuce/Leafy Greens Commodity Specific Guidance**  
273 **Production & Harvest Unit Operations**  
274

275 **1. PURPOSE**

276 The issues identified in this document are based on the core elements of Good Agricultural  
277 Practices. The specific recommendations contained herein are intended for lettuce and leafy  
278 greens only. If these specific recommendations are effectively implemented this would  
279 constitute the best practices for a GAP program for the production and harvest unit operations  
280 of lettuce and leafy greens.  
281

282 **2. ISSUE: GENERAL REQUIREMENTS**

283 In addition to the area-specific requirements discussed in latter sections, there are several  
284 general requirements that are part of an effective best practices program. These requirements  
285 are outlined below.  
286

287 **2.1. The Best Practices Are:**

- 288 • A written Leafy Greens Compliance Plan which specifically addresses the Best  
289 Practices of this document shall be prepared. This plan shall address at least the  
290 following areas: water, soil amendmets, environmental factors, work practices,  
291 and field sanitation.
- 292 • Shippers shall have an up to date producers list with contact and location  
293 information on file.
- 294 • The shipper shall comply with the requirements of The Public Health Security  
295 and Bioterrorism Preparedness and Response Act of 2002 (farms are exempt  
296 from the Act) including those requirements for recordkeeping (traceability) and  
297 registration.
- 298 • Each producer and shipper shall designate an individual responsible for their  
299 operation's food safety program. Twenty-four hour contact information shall be  
300 available for this individual in case of food safety emergencies.

301  
302 **3. ISSUE: ENVIRONMENTAL ASSESSMENTS**

303 This section addresses assessments that shall be completed prior to the first seasonal planting,  
304 within one week prior to harvesting and during harvest operations. These environmental  
305 assessments are intended to identify any issues related to the produce field, adjacent land  
306 uses, or intrusion by animal of significant risk (see Table 5) that might impact produce safety.  
307

308 **3.1. The Best Practices Are:**

- 309 • Prior to the first seasonal planting and within one week prior to harvest, perform  
310 an environmental assessment of the production field and surrounding area. Focus

311 these assessments on evaluating the production field for possible animal of  
312 significant risk intrusion or other sources of human pathogens of concern,  
313 assessing adjacent land uses for possible sources that might contaminate the  
314 production field, and evaluating nearby water sources for the potential of past or  
315 present flooding.

- 316 ○ Assessment of Produce Field
  - 317 ■ Evaluate all produce fields for evidence of animal of significant
  - 318 risk intrusion and/or feces. If any evidence is found, follow
  - 319 procedures identified in the “Production Locations -
  - 320 Encroachment by Animals and Urban Settings.”
- 321 ○ Assessment of Adjacent Land Use
  - 322 ■ Evaluate all land and waterways adjacent to all production fields
  - 323 for possible sources of human pathogen of concern. These
  - 324 sources include, but are not limited to, manure storage, compost
  - 325 storage, CAFO’s, grazing/open range areas, surface water,
  - 326 sanitary facilities, and composting operations (see Table 6 for
  - 327 further detail). If any possible uses that might result in produce
  - 328 contamination are present, follow management practices identified
  - 329 in the sections below related to environmental and land use
  - 330 concerns.
- 331 ○ Assessment of Historical Land Use
  - 332 ■ To the degree practical, determine and document the historical
  - 333 land uses for production fields and any potential issues from these
  - 334 uses that might impact food safety (i.e., hazardous waste sites,
  - 335 landfills, etc.).
- 336 ○ Assessment of Flooding
  - 337 ■ Evaluate all produce fields for evidence of flooding. If any
  - 338 evidence is found, follow procedures identified in the “Flooding”
  - 339 section below.
  - 340

341 **4. ISSUE: WATER**

342 Water used for production and harvest operations may contaminate lettuce and leafy greens if  
343 water containing human pathogens comes in direct contact with the edible portions of  
344 lettuce/leafy greens. Contamination may also occur by means of water-to-soil followed by  
345 soil-to-lettuce/leafy greens contact. Irrigation methods may have varying potential to  
346 introduce human pathogens or promote human pathogen growth on lettuce and leafy greens  
347 (Stine *et al.*, 2005).

348  
349 There are several different approaches and values that can be utilized to ensure that water is  
350 of appropriate quality for its intended use. The metrics applied in this edition of the  
351 Commodity Specific Guidance should be considered a starting point in industry efforts to  
352 continuously improve the quality of water used in production of these commodities.  
353

354 The current metrics are intended to provide standards associated with water uses; however, it  
355 is known that various water sources have different microbial qualities, and each source  
356 should be monitored accordingly. Typical microbial values associated with various sources

357 can be found in the Sanitary Survey document (Appendix A). During the sanitary survey that  
358 is performed prior to each growing season expected microbial values and historical  
359 monitoring data should be used to evaluate the quality of the water source.  
360

361 **4.1. The Best Practices Are:**

- 362 • A water system description shall be prepared. This description can use maps,  
363 photographs, drawings or other means to communicate the location of permanent  
364 fixtures and the flow of the water system (including any water captured for re-  
365 use.). Permanent fixtures include wells, gates, reservoirs, valves, returns and  
366 other above ground features that make up a complete irrigation system should be  
367 documented in such a manner as to enable location in the field. Water sources  
368 and the production blocks they may serve should be documented.
- 369 • Water systems that convey untreated human or animal waste must be separated  
370 from conveyances utilized to deliver irrigation water.
- 371 • Use irrigation water and water in harvest operations that is of appropriate  
372 microbial quality for its intended use; see Table 1 and Decision Trees (1A, 1B  
373 and 1C) for specific numerical criteria. Appendix B provides the basis for these  
374 water quality metrics.
- 375 • Perform a sanitary survey prior to use of water in agricultural operations and if  
376 water quality microbial tests are at levels that exceed the numerical values set  
377 forth in Table 1. The sanitary survey is described in Appendix A.
- 378 • Test water as close to the point-of-use as practical, and if microbial levels are  
379 above specific action levels, take appropriate remedial and corrective actions.
- 380 • Retain documentation of all test results and/or Certificates of Analysis available  
381 for inspection for a period of at least 2 years.

382 Other Considerations for water

- 383 ○ Evaluate irrigation methods (drip irrigation, overhead sprinkler, furrow, etc.)  
384 for their potential to introduce, support or promote the growth of human  
385 pathogens on lettuce and leafy greens. Consider such factors as the potential  
386 for depositing soil on the crop, presence of pooled or standing water that  
387 attracts animals, etc.
- 388 ○ When waters from various sources are combined, consider the potential for  
389 pathogen growth in the water.
- 390 ○ For surface water sources, consider the impact of storm events on irrigation  
391 practices. Bacterial loads in surface water are generally much higher after a  
392 storm than normal, and caution shall be exercised when using these waters for  
393 irrigation.
- 394 ○ Use procedures for storing irrigation pipes and drip tape that reduce or  
395 eliminate potential pest infestations. Develop procedures to provide for  
396 microbiologically safe use of irrigation pipes and drip tape if a pest  
397 infestation does occur.

398                   ○ Reclaimed water shall be subject to applicable state and federal regulations  
399                   and standards. Use of this water for agricultural purposes must meet the most  
400                   stringent standard as defined by the following: state and federal regulation or  
401                   Table 1 of this document.

402   **5.        ISSUE: WATER USAGE TO PREVENT PRODUCT DEHYDRATION**

403   Lettuce/leafy greens may be sprayed with small amounts of water during machine harvest or  
404   in the field container just after harvest to reduce water loss. Water used in harvest operations  
405   may contaminate lettuce and leafy greens if there is direct contact of water containing human  
406   pathogens with edible portions of lettuce/leafy greens.  
407

408       **5.1.        The Best Practices Are:**

- 409           • Due to the timing of application of water that directly contacts edible portions of  
410           lettuce/leafy greens, assure the water is of appropriate microbial quality (e.g.,  
411           meets U.S. EPA microbial standards for drinking water).
- 412           • Test the water source periodically to demonstrate it is of appropriate microbial  
413           quality for its intended purpose (e.g., meets U.S. EPA or WHO microbial  
414           standards for drinking water) or assure that it has appropriate disinfection  
415           potential as described in Table 1.

416 TABLE 1. WATER USE

Use	Metric	Rationale /Remedial Actions
<p><b>PREHARVEST Foliar Applications</b> Whereby Edible Portions of the Crop <b>ARE</b> Contacted by Water</p> <p>(e.g. overhead sprinkler irrigation, pesticides/fungicide application, etc.)</p>	<p><b>Target Organism:</b> generic <i>E. coli</i>.</p> <p><b>Sampling Procedure:</b> 100 mL sample collected aseptically at the point of use; i.e., one sprinkler head per water source for irrigation, water tap for pesticides, etc. Water utilized in preseason irrigation operations may be tested and utilized.</p> <p><b>Sampling Frequency:</b> One sample per water source shall be collected and tested prior to use if &gt;60 days since last test of the water source. Additional samples shall be collected no less than 18 hr apart and at least monthly during use from points within the distribution system.</p> <p><b>Municipal &amp; Well Exemption:</b> For wells and municipal water sources, if generic <i>E. coli</i> are below detection limits for five consecutive samples, the sampling frequency may be decreased to no less than once every 180 days and the requirements for 60 and monthly sampling are waived. This exemption is void if there is a significant source or distribution system change.</p>	<p>For any given water source (municipal, well, reclaimed water, reservoir or other surface water), samples for microbial testing shall be taken at a point as close to the point of use as practical (as determined by the sampler, to ensure the integrity of the sample, using sampling methods as prescribed in Table 1) where the water contacts the crop, so as to test both the water source and the water distribution system. Only one sample per month per distribution system is required under these metrics. If there are multiple potential point-of-use sampling points in a distribution system, then samples shall be taken from different point-of-use locations each subsequent month (randomize or rotate sample locations).</p> <p>Water for preharvest, direct edible portion contact shall meet or exceed microbial standards for recreational water, based on a rolling geometric mean of the five most recent samples. <b>However, a rolling geometric mean of five samples is not necessarily required prior to irrigation or harvest. If less than five samples are collected prior to irrigation, the acceptance criteria depends on the number of samples taken. If only one sample has been taken, it must be below 126 CFU/100 mL. Once two samples are taken, a geometric mean can be calculated and the normal acceptance criteria apply. If the acceptance criteria are exceeded during this time period, additional samples may be collected to reach a 5 sample rolling geometric mean (as long as the water has not been used for irrigation). The rolling geometric mean calculation starts after 5 samples have been collected.</b> If the water source has not been tested in the past 60 days, the first water sample shall be tested prior to use, to avoid using a contaminated water source. After the first sample is shown to be within acceptance criteria, subsequent samples shall be collected no less frequently than monthly at points of use within the distribution system.</p> <p>Ideally, preharvest water should not contain generic <i>E. coli</i>, but low levels do not necessarily indicate that the water is unsafe. Investigation and/or remedial action <b>SHOULD</b> be taken when test results are higher than normal, or indicate an upward trend. Investigation and remedial action <b>SHALL</b> be taken when acceptance criteria are exceeded.</p> <p><b>Remedial Actions:</b> If the rolling geometric mean (n=5) or any one sample exceeds the acceptance criteria, then the water shall not be used whereby edible portions of the crop are contacted by water until remedial actions have been completed and generic <i>E. coli</i> levels are within acceptance criteria:</p> <ul style="list-style-type: none"> <li>• Conduct a sanitary survey of water source and distribution system to determine if a contamination source is evident and can be eliminated. Eliminate identified contamination source(s).</li> <li>• For wells, perform a sanitary survey and/or treat as described in Appendix A Sanitary Survey.</li> <li>• Retest the water after conducting the sanitary survey and/or taking remedial actions to determine if it meets the outlined microbial acceptance criteria for this use. This sample should represent the</li> </ul>

	<p><b>Test Method:</b> 15 tube MPN (FDA BAM) or other U.S. EPA, AOAC, or other method accredited for quantitative monitoring of water for generic <i>E. coli</i>. Presence/absence testing with a similar limit of detection may be used as well.</p> <p><b>Acceptance Criteria:</b> ≤126 MPN (or CFU*)/100 mL (rolling geometric mean n=5) and ≤235 MPN/100mL for any single sample.</p> <p>*for the purposes of water testing, MPN and CFU shall be considered equivalent.</p>	<p>conditions of the original water system, if feasible this test should be as close as practical to the original sampling point. A more aggressive sampling program (i.e., sampling once per week instead of once per month) shall be instituted if an explanation for the exceedence is not readily apparent. This type of sampling program should also be instituted if an upward trend is noted in normal sampling results.</p> <p><b>Crop Testing:</b> If water testing indicates that a crop has been directly contacted with water exceeding acceptance criteria, product shall be sampled and tested for <i>E. coli</i> O157:H7 and <i>Salmonella</i> as described in Appendix C, prior to harvest. If crop testing indicates the presence of either pathogen, the crop shall NOT be harvested for human consumption.</p> <p><b>Records:</b> All test results and remedial actions shall be documented and available for verification from the producer who is the responsible party for a period of two years.</p>
<p><b>PREHARVEST Non-foliar Applications</b> Whereby Edible Portions of the Crop are <b>NOT</b> Contacted by Water</p> <p>(e.g., furrow or drip irrigation, dust abatement water; if water is not used in the vicinity of produce, then testing is not necessary)</p>	<p><b>Target Organism, Sampling Procedure, Sampling Frequency Test Method and Municipal Well Exemption:</b> as described for foliar application.</p> <p><b>Acceptance Criteria:</b> ≤126 MPN /100 mL (rolling geometric mean n=5) and ≤576 MPN /100 mL for any single sample.</p>	<p>Testing and remedial actions for preharvest water that does not come in direct contact with edible portions of the crop are the same as for direct contact water, but acceptance criteria are less stringent because of the reduced risk of contact of the edible portion with contamination from water. Acceptance criteria here are derived from U.S. EPA recreational water standards.</p>

<p><b>POSTHARVEST Direct Product Contact or Food Contact Surfaces</b></p>	<p><b><u>Microbial Testing</u></b>  <b>Target Organism, Sampling Procedure, and Test Method:</b> as described for foliar application.</p> <p><b>Sampling Frequency:</b> One sample per water source shall be collected and tested prior to use if &gt;60 days since last test of the water source. Additional samples shall be collected at intervals of no less than 18 hr and at least monthly during use.</p> <p><b>Acceptance Criteria:</b>  <b>Negative or below DL for all samples</b></p>	<p>Water that directly contacts edible portions of harvested crop, or is used on food contact surfaces, such as equipment or utensils, shall meet the Maximum Contaminant Level Goal for <i>E. coli</i> as specified by U.S. EPA or contain an approved disinfectant at sufficient concentration to prevent cross contamination. Microbial or physical/chemical testing shall be performed, as appropriate to the specific operation, to demonstrate that acceptance criteria have been met.</p> <p><b>Single Pass vs. Multiple Pass Systems</b></p> <ul style="list-style-type: none"> <li>• Single pass use – Water must have non-detectable levels of <i>E. coli</i> or breakpoint disinfectant present at point of entry</li> <li>• Multi-pass use – Water must have non-detectable levels of <i>E. coli</i> and/or sufficient disinfectant to insure returned water has no detectable <i>E. coli</i> (minimally 1 ppm chlorine)</li> </ul> <p><b>Remedial Actions:</b>  If any one sample exceeds the acceptance criteria, then the water shall not be used for this purpose unless appropriate disinfectants have been added or until remedial actions have been completed and generic <i>E. coli</i> levels are within acceptance criteria:</p> <ul style="list-style-type: none"> <li>• Conduct a sanitary survey of water source and distribution system to determine if a contamination source is evident and can be eliminated. Eliminate identified contamination source(s).</li> <li>• For wells, perform a sanitary survey and/or treat as described in Appendix A Sanitary Survey.</li> <li>• Retest the water at the same sampling point after conducting the sanitary survey and/or taking remedial actions to determine if it meets the outlined microbial acceptance criteria for this use.</li> </ul> <p>For example, if a water sample for water used to clean food contact surfaces has detectable <i>E. coli</i>, STOP using that water system, examine the distribution line and source inlet as described in Appendix A Sanitary Survey, and retest from the same point of use. Continue testing daily for 5 days at the point closest to use, and do not use the water system until it consistently delivers water that is safe, sanitary water and of appropriate microbial quality (i.e. Negative result) for the intended use. If any of the any of the five samples taken during the intensive sampling period after corrective actions have been taken have detectable <i>E. coli</i>, repeat remedial actions and DO NOT use that system until the source of contamination can be corrected.</p> <p><b>Records:</b> All test results and remedial actions shall be documented and available for verification from the user of the water for a period of two years.</p>
	<p><b><u>Physical/Chemical Testing</u></b>  <b>Target Variable:</b>  Water disinfectant (e.g. chlorine or other disinfectant compound, ORP)</p> <p><b>Multi Pass Water Acceptance Criteria:</b></p> <ul style="list-style-type: none"> <li>• <u>Chlorine</u>  ≥1 ppm free chlorine after application and pH 6.5 – 7.5 OR</li> <li>• ORP ≥ 650 mV, and pH 6.5 – 7.5</li> <li>• <u>Other approved treatments</u> per product EPA label for human pathogen reduction in water.</li> </ul> <p><b>Testing Procedure:</b></p> <ul style="list-style-type: none"> <li>• Chemical reaction based colorimetric test, or</li> <li>• Ion specific probe, or</li> <li>• ORP, or</li> <li>• Other as recommended by disinfectant supplier.</li> </ul>	


	<p><b>Testing Frequency:</b> Continuous monitoring (preferred) with periodic verification by titration OR Routine monitoring if the system can be shown to have a low degree of variation.</p>	
--	--	--

417 **Figure 1A. Decision Tree for PRE-HARVEST WATER USE – Foliar Applications**  
 418 **whereby edible portions of the crop are contacted by water (e.g. overhead irrigation,**  
 419 **pesticide/fungicide applications)**

For any given water source (municipal, well, reclaimed water, reservoir or other surface water):

**Sampling Frequency:** One sample per water source shall be collected and tested prior to use if >60 days since last test of the water source. Additional samples shall be collected at intervals of no less than 18 hr and at least monthly during use.

- Sample sources as close to the point-of-use as practical, as determined by the sampler to ensure the integrity of the sample, using sampling methods as prescribed in Table 1.
- Analyze samples for generic *E. coli* using a 15-tube MPN methodology. Other EPA-, FDA- or AOAC- or other accredited method may be used.
- Geometric means, including rolling geometric means shall be calculated using the five most recent samples.



**Acceptance Criteria**  
 ≤ 126 MPN/100ml  
 (geometric mean of five samples)  
**AND**  
 ≤235 MPN/100ml (all single samples)

**Action Level**  
 > 126 MPN/100ml  
 (geometric mean of five samples)  
**OR**  
 >235 MPN/100ml (any single sample)

No further action necessary. Water from this source may be used for any pre-harvest use such as crop foliar applications and/or irrigation.

However, when test results are higher than normal or indicate an upward trend, investigation and/or remedial action SHOULD be taken.

**Remedial Actions:**


- Discontinue use for foliar and direct contact with the edible portion of the plant applications until it returns to compliance.
- Examine the water source and distribution system to determine if a contamination source is evident and can be eliminated.
- For wells, perform a sanitary survey and/or treat as described in Appendix A Sanitary Survey.
- After sanitary survey and/or remedial actions have been taken, retest the water at the same sampling point.
- Test daily for five days, approximately 24h apart, at the point closest to use.
- If any of the next five samples is >235 MPN/100mL, repeat sanitary survey and/or remedial action.
- Do not use water from that water system, in a manner that directly contact edible portions of the crop, until the water can meet the outlined acceptance criteria for this use.

**Crop testing:**

- If crop has been directly contacted with water exceeding acceptance criteria, sample and test product for *E. coli* O157:H7 and *Salmonella* as described in Appendix C, prior to harvest.
- If crop testing indicates the presence of either pathogen, do NOT harvest for human consumption.

421  
422  
423  
424


**Figure 1B. Decision Tree for PRE-HARVEST WATER USE – Non-Foliar Applications whereby edible portions of the crop are NOT contacted by water (e.g. furrow or drip irrigation, dust abatement water)**



425

426  
427

**Figure 1C. POSTHARVEST WATER USE – Direct product contact (e.g. re-hydration, core in field, etc.)**



428  
429  
430

431 **6. ISSUE: SOIL AMENDMENTS**

432 Soil amendments are commonly but not always incorporated prior to planting into  
433 agricultural soils used for lettuce/leafy greens production to add organic and inorganic  
434 nutrients to the soil as well as intended to improve the physical, chemical, or biological  
435 characteristics of soil.. Human pathogens may persist in animal manures for weeks or even  
436 months (Fukushima *et al.* 1999; Gagliardi and Karns 2000). Proper composting of animal  
437 manures via thermal treatment will reduce the risk of potential human pathogen survival.  
438 However, the persistence of many human pathogens in agricultural soils depends on many  
439 factors (soil type, relative humidity, UV index, etc.) and the effects of these factors is under  
440 extensive investigation (Jiang *et al.* 2003; Islam *et al.* 2004).

441  
442 Field soil contaminated with human pathogens may provide a means of lettuce and leafy  
443 greens contamination. Studies of human pathogens conducted in cultivated field vegetable  
444 production models point towards a rapid initial die-off from high pathogen populations but a  
445 characteristic and prolonged low level survival. Readily detectable survival is typically less  
446 than 8 weeks following incorporation, but has been documented to exceed 12 weeks (Jiang *et*  
447 *al.* 2001; Islam *et al.* 2005).. Recoverable pathogen populations, using highly sensitive  
448 techniques, have been reported to persist beyond this period under some test conditions. The  
449 detection of introduced pathogens on mature lettuce plants from these low levels of surviving  
450 pathogens was not possible, and the risk was concluded to be negligible. Human pathogens  
451 do not persist for long periods of time in high UV index and low relative humidity  
452 conditions, but may persist for longer periods of time within aged manure or inadequately  
453 composted soil amendments. Therefore, establishing suitably conservative pre-plant  
454 intervals, appropriate for specific regional and field conditions, is an effective step towards  
455 minimizing risk (Suslow *et al.* 2003).

456

457 **6.1. The Best Practices Are:**

- 458 • DO NOT USE raw manure or soil amendment that contain un-composted,  
459 incompletely composted or non-thermally treated animal manure to fields which  
460 will be used for lettuce and leafy green production.
- 461 • See Table 2 and Decision Trees (Figures 2A and 2B) for numerical criteria and  
462 guidance for compost and soil amendments used in lettuce and leafy greens  
463 production fields. The “Technical Basis Document” (Appendix B) describes the  
464 process used to develop these metrics.
- 465 • Any soil amendment that does not contain animal manure must have a document  
466 (e.g., ingredient list, statement of identity, letter of guaranty, etc.) from the  
467 producer or seller demonstrating that it is manure free. This document must  
468 indicate in some way that manure is not an ingredient used in the production of  
469 the amendment or provide the ingredients of the product. A statement of identity  
470 or product is sufficient for single-chemical amendments (i.e., “calcium  
471 carbonate” or “gypsum”). If “inert ingredients” are listed as part of an  
472 amendment, then a document from the producer or seller is necessary indicating  
473 manure has not been added. The manure free document must be available for  
474 verification before harvest begins and it must be saved and available for  
475 inspection for 2 years. A new document is required every two years unless there  
476 is a significant process or ingredient change.

- 477 • Implement management plans (e.g., timing of applications, storage location,  
478 source and quality, transport, etc.) that significantly reduce the likelihood that soil  
479 amendments being used contain human pathogens.
- 480 • Verify that the time and temperature process used during the composting process  
481 reduces, controls, or eliminates the potential for human pathogens being carried  
482 in the composted materials, as applicable to regulatory requirements.
- 483 • Maximize the time interval between soil amendment application and time to  
484 harvest.
- 485 • Implement practices that control, reduce or eliminate likely contamination of  
486 lettuce/leafy green fields in close proximity to on-farm stacking of manure.
- 487 • Use soil amendment application techniques that control, reduce or eliminate  
488 likely contamination of surface water and/or edible crops being grown in adjacent  
489 fields.
- 490 • Segregate equipment used for soil amendment handling, preparation, distribution,  
491 applications or use effective means of equipment sanitation before subsequent use  
492 that effectively reduce the potential for cross contamination.
- 493 • Minimize the proximity of wind-dispersed or aerosolized sources of  
494 contamination (e.g., water and manure piles) that may potentially contact growing  
495 lettuce/leafy greens or adjacent edible crops. Segregate equipment used for soil  
496 amendment applications or use effective means of equipment sanitation before  
497 subsequent use.
- 498 • Compost suppliers shall have written Standard Operating Procedures to prevent  
499 cross-contamination of finished compost with raw materials through equipment,  
500 runoff, or wind, and producers shall obtain proof that these documents exist.
- 501 • Compost operations supplying compost to leafy greens crops shall maintain  
502 temperature monitoring and turning records for at least two years, and producers  
503 shall obtain proof that this documentation exists. This applies to composting  
504 operations regulated under Title 14 CCR as well as smaller operations that do not  
505 fall under Title 14.
- 506 • Perform microbiological testing of soil amendments prior to application (Table  
507 2).
- 508 • Do not use biosolids as a soil amendment for production of lettuce or leafy  
509 greens.
- 510 • Retain documentation of all processes and test results by lot (at the supplier)  
511 and/or Certificates of Analysis available for inspection for a period of at least two  
512 years.
- 513
- 514

TABLE 2. SOIL AMENDMENTS

Amendment	Metric/Rationale
<p><b>Raw Manure or Not Fully Composted Animal Manure Containing Soil Amendments (see composted manure process definition below)</b></p>	<p><b>DO NOT USE OR APPLY</b> soil amendments that contain un-composted, incompletely composted or non-thermally treated (e.g., heated) animal manure to fields which will be used for lettuce and leafy greens production. If these materials have been applied to a field, wait one year prior to producing leafy greens.</p>
<p><b>Composted Soil Amendments (containing animal manure or animal products)</b></p> <p>*Composted soil amendments should not be applied after emergence of plants.</p>	<p><b>Please see Figure 2A: Decision Tree for Use of Composted Soil Amendments.</b></p> <p><b>Composting Process Validation:</b></p> <p><u>Enclosed or within-vessel composting:</u> Active compost must maintain a minimum of 131°F for 3 days</p> <p><u>Windrow composting:</u> Active compost must maintain aerobic conditions for a minimum of 131°F for 15 days, with a minimum of five turnings.</p> <p><u>Aerated static pile composting:</u> Active compost must be covered with at least 12 inches of insulating materials and maintain a minimum of 131°F for 3 days</p> <p><b>Target Organisms:</b></p> <ul style="list-style-type: none"> <li>• Fecal coliforms</li> <li>• <i>Salmonella</i> spp</li> <li>• <i>E. coli</i> O157:H7</li> </ul> <p><b>Acceptance Criteria:</b></p> <ul style="list-style-type: none"> <li>• Fecal coliforms &lt;1000 MPN/gram</li> <li>• <i>Salmonella</i>: Negative or &lt; DL (&lt;1/ 30 grams)</li> <li>• <i>E. coli</i> O157:H7: Negative or &lt; DL (&lt;1/ 30 grams)</li> </ul>

Amendment	Metric/Rationale
	<p><b>Recommended Test Methods:</b></p> <ul style="list-style-type: none"> <li>• Fecal coliforms: 9 tube MPN</li> <li>• <i>Salmonella spp.</i>: U.S. EPA Method 1682</li> <li>• <i>E. coli</i> O157:H7: Any laboratory validated method for compost sampling.</li> <li>• Other U.S. EPA, FDA, or AOAC-accredited methods may be used as appropriate.</li> </ul> <p><b>Sampling Plan:</b></p> <ul style="list-style-type: none"> <li>• 12 point sampling plan composite sample (divide each lot/pile into a 3 x 4 grid and extract 12 equivolume samples.)</li> <li>• Sample may be taken by the supplier if trained by the testing laboratory</li> <li>• Laboratory must be certified/accredited for microbial testing by an appropriate process authority</li> </ul> <p><b>Testing Frequency:</b></p> <ul style="list-style-type: none"> <li>• Each lot before application to production fields. A lot is defined as a unit of production equal to or less than 5,000 cubic yards.</li> </ul> <p><b>Application Interval:</b></p> <ul style="list-style-type: none"> <li>• Must be applied &gt;45 days before harvest</li> </ul> <p><b>Documentation:</b></p> <ul style="list-style-type: none"> <li>• All test results and/or Certificates of Analysis shall be documented and available for verification from the producer (the responsible party) for a period of two years.</li> </ul> <p><b>Rationale:</b></p> <ul style="list-style-type: none"> <li>• The microbial metrics and validated processes for compost are based on allowable levels from California state regulations (CCR Title 14 - Chapter 3.1 - Article 5 2007), with the addition of testing for <i>E. coli</i> O157:H7 as microbe of particular concern. The 45-day application interval was deemed appropriate due to the specified multiple hurdle risk reduction approach outlined. Raw manure must be composted with an approved process and pass testing requirements before an application.</li> </ul>

<p><b>Soil amendments containing animal manure that has been physically heat treated or processed by other equivalent methods.</b></p>	<p><b>Please see Figure 2B: Decision Tree for Use of Physically Heat Treated Soil Amendments.</b></p> <p><b>Physical Heat Process Validation</b></p> <ul style="list-style-type: none"> <li>The physical heat treatment processes applied to the soil amendment containing animal manure shall be done via a process validated to assure that the process is capable of reducing pathogens of human health significance to acceptable levels.</li> </ul> <p><b>Target Organism:</b></p> <ul style="list-style-type: none"> <li>Fecal coliforms</li> <li><i>Salmonella</i> spp</li> <li><i>E. coli</i> O157:H7</li> </ul> <p><b>Acceptance Criteria:</b></p> <ul style="list-style-type: none"> <li>Fecal coliforms Negative or &lt; DL per gram</li> <li><i>Salmonella</i>: Negative or &lt; DL (&lt;1/ 30 grams)</li> <li><i>E. coli</i> O157:H7: Negative or &lt; DL (&lt;1/ 30 grams)</li> </ul> <p><b>Recommended Test Methods:</b></p> <ul style="list-style-type: none"> <li>Fecal coliforms: 9 tube MPN</li> <li><i>Salmonella</i> spp: U.S. EPA Method 1682</li> <li><i>E. coli</i> O157:H7: Any laboratory validated method for testing soil amendments.</li> <li>U.S. EPA, FDA, AOAC-or other accredited methods may be used as appropriate</li> </ul> <p><b>Sampling Plan:</b></p> <ul style="list-style-type: none"> <li>12 point sampling plan composite sample (divide each lot/pile into a 3 x 4 grid and extract 12 equivolume samples)</li> <li>Sample may be taken by the supplier if trained by the testing laboratory</li> <li>Laboratory must be certified/accredited by annual review of laboratory protocols based on GLPs by recognized NGO.</li> </ul> <p><b>Testing Frequency:</b></p> <ul style="list-style-type: none"> <li>Each lot before application to production fields. <ul style="list-style-type: none"> <li>In lieu of the above analysis requirement a Certificate of Process Validity Issued by a recognized <i>Process Authority</i> can be substituted. This certificate will attest to the process validity as determined by either a documented (included w/Certificate)) inoculated pack study</li> </ul> </li> </ul>
--	---

of the standard process or microbial inactivation calculations of organisms of significant risk (included w/Certificate) as outlined in FDA CFSAN publication “**Kinetics of Microbial Inactivation for Alternative Food Processing Technologies. Overarching Principles: Kinetics and Pathogens of Concern for All Technologies**” (Incorporated for reference in Appendix E Thermal Process Overview)

**Application Interval:**

- If the physical heat treatment process used to inactivate human pathogens of significant public health concern that may be found in animal manure containing soil amendments, is validated and meets the microbial acceptance criteria outlined below, then no time interval is needed between application and harvest.
- If the physical heat treatment process used to inactivate human pathogens of significant public health concern that may be found in animal manure containing soil amendments is not validated but will likely significantly reduce microbial populations of human pathogens (minimum temperature: 300°F (150°C) for 60 minutes resulting in a moisture content <30% dry weight) and meets that microbial acceptance criteria outlined above, then a 45 day interval between application and harvest is required.

**Documentation:**

- All test results and/or Certificates of Analysis and/or Certificates of Process Validation shall be documented and available for verification from the producer who is the responsible party for a period of two years. The suppliers operation should be validated by a process authority and a record maintained by the producer for a period of two years.

**Rationale:**


- The microbial metrics and validated processes for compost are based on allowable levels from California state regulations (CCR Title 14 - Chapter 3.1 - Article 5 2007), with the addition of testing for *E. coli* O157:H7 as the microbe of particular concern. A more stringent level of fecal coliform was also included to address the much more controlled nature of soil amendments produced in this manner. The above suggested application interval was deemed appropriate due to the specified multiple hurdle risk reduction approach outlined. Raw manure must be composted with an approved process and pass testing requirements before application.
- FDA has established the validity of D-values and Z-values for key pathogens of concern in foods. This method of process validation is currently acceptable to US regulators. Alternatively, results of an inoculated test pack utilizing the specific process is also an acceptable validation of the lethality of the process.

<b>Soil Amendments Not Containing Animal Manure</b>	<ul style="list-style-type: none"><li>• Any soil amendment that DOES NOT contain animal manure must have documentation that it is manure-free.</li><li>• The documentation must be available for verification before harvest begins.</li><li>• If there is documentation that the amendment does not contain manure or animal products then no additional testing is required, and there is no application interval necessary</li><li>• Any test results and/or documentation shall be available for verification from the producer who is the responsible party for a period of two years.</li></ul>
---	---

517  
518  
519

**Figure 2A. Decision Tree for Composted Soil Amendments (SA)**


If raw manure has been directly applied to the field in the past, a 1 year waiting period shall be observed before planting any variety of leafy green crops.



520

521  
522  
523

**Figure 2B. Decision Tree for Physically Heat Treated Animal Manure Containing Soil Amendments (SA)**



524  
525

526 **7. ISSUE: NONSYNTHETIC CROP TREATMENTS**

527 Nonsynthetic crop treatments are commonly applied post-emergence for pest and disease  
528 control, greening, and to provide organic and inorganic nutrients to the plant during the  
529 growth cycle. For the purposes of this document, they are defined as any crop input that  
530 contains animal manure, an animal product, and/or an animal by-product that is reasonably  
531 likely to contain human pathogens. Due to the potential for human pathogen contamination,  
532 these treatments should only be used under conditions that minimize the risk for crop  
533 contamination.  
534

535 **7.1. The Best Practices Are:**

- 536 • Do not use crop treatments that contain raw manure for lettuce or leafy green  
537 produce.
- 538 • Retain documentation of all test results available for inspection for a period of at  
539 least two years.
- 540 • Implement management plans (e.g. timing of applications, storage location,  
541 source and quality, transport, etc.) that assure to the greatest degree practicable  
542 that the use of crop treatments does not pose a significant pathogen contamination  
543 hazard.
- 544 • Verify that the time and temperature process used during crop treatment  
545 manufacture reduces, controls, or eliminates the potential for human pathogens  
546 being carried in the composted materials, as applicable to regulatory  
547 requirements.
- 548 • Maximize the time interval between the crop treatment application and time to  
549 harvest.
- 550 • Implement practices that control, reduce or eliminate likely contamination of  
551 lettuce/leafy green fields that may be in close proximity to on-farm storage of  
552 crop treatments.
- 553 • Use crop treatment application techniques that control, reduce or eliminate the  
554 likely contamination of surface water and/or edible crops being grown in adjacent  
555 fields.
- 556 • Segregate equipment used for crop treatment applications or use effective means  
557 of equipment sanitation before subsequent use.
- 558 • See Table 3 and Decision Tree (Figure 3) for numerical criteria and guidance for  
559 nonsynthetic crop treatments used in lettuce and leafy greens production fields.  
560 The “Technical Basis Document” (Appendix B) describes the process used to  
561 develop these metrics.

562

563

564

TABLE 3. NONSYNTHETIC CROP TREATMENTS


Treatment	Metric/Rationale
<p><i>Any crop input that contains animal manure, an animal product, and/or an animal by-product that is reasonably likely to contain human pathogens.</i></p> <p>Examples include but are not limited to:</p> <ul style="list-style-type: none"> <li>• Compost teas,</li> <li>• Fish emulsions</li> <li>• Fish meal</li> <li>• Blood meal</li> <li>• "Bio-fertilizers" commonly used for pest control, greening, disease control, fertilizing.</li> </ul> <p>Suppliers of these products shall disclose on labels, certificates of analysis, or other companion paperwork whether the product contains any animal manure or products.</p>	<p><b>Non synthetic crop treatments that contain animal products or animal manure that have not been physically heat treated or processed by other equivalent methods shall NOT be directly applied to the edible portions of lettuce and leafy greens.</b></p> <p><b>Please see Figure 3: Decision Tree for Use of Nonsynthetic Crop Treatments.</b></p> <p><b>Process Validation</b></p> <ul style="list-style-type: none"> <li>• The physical, chemical and/or biological treatment process(es) used to render the crop input safe for application to edible crops must be validated.</li> </ul> <p><b>Target Organism:</b></p> <ul style="list-style-type: none"> <li>• <i>Salmonella</i> spp</li> <li>• <i>E. coli</i> O157:H7</li> </ul> <p><b>Acceptance Criteria (at point of use):</b></p> <ul style="list-style-type: none"> <li>• <i>Salmonella</i>: Negative or &lt; DL (&lt;1/ 30 grams)</li> <li>• <i>E. coli</i> O157:H7: Negative or &lt; DL (&lt;1/ 30 grams)</li> <li>• Other pathogens appropriate for the source material</li> </ul> <p><b>Recommended Test Methods:</b></p> <ul style="list-style-type: none"> <li>• <i>Salmonella</i> spp: U.S. EPA Method 1682</li> <li>• <i>E. coli</i> O157:H7: Any laboratory validated method for the non synthetic material to be tested.</li> <li>• Other U.S. EPA, FDA, or AOAC-accredited methods may be used as appropriate</li> </ul> <p><b>Sampling Plan:</b></p> <ul style="list-style-type: none"> <li>• 12 point sampling plan composite sample (if solid), one sample per batch if liquid (if liquid-based, then water quality acceptance levels as described in Table 1 should be used)</li> <li>• Sample may be taken by the supplier if trained by the testing laboratory</li> <li>• Laboratory must be certified/accredited by annual review of laboratory protocols based on GLPs by recognized NGO</li> </ul> <p><b>Testing Frequency:</b></p> <ul style="list-style-type: none"> <li>• Each lot before application to production fields.</li> </ul>

Treatment	Metric/Rationale
	<p><b>Application Interval:</b></p> <ul style="list-style-type: none"> <li>• If the physical, chemical and/or biological treatment process used to render the crop input safe for application to edible crops is validated and meets that microbial acceptance criteria outlined above, no time interval is needed between application and harvest.</li> <li>• If the physical, chemical and/or biological treatment process used to render the crop input safe for application to edible crops is not validated yet meets the microbial acceptance criteria outlined above, a 45 day time interval between application and harvest is required.</li> </ul> <p><b>Documentation:</b></p> <ul style="list-style-type: none"> <li>• All test results and/or Certificates of Analysis shall be documented and available from the producer for verification for a period of 2 years. The producer the party responsible party for maintaining the appropriate records.</li> </ul> <p><b>Rationale:</b></p> <ul style="list-style-type: none"> <li>• The microbial metrics and validated processes for compost are based on allowable levels from California state regulations (CCR Title 14 - Chapter 3.1 - Article 5 2007), with the addition of testing for <i>E. coli</i> O157:H7 as the microbe of particular concern. The above suggested application interval was deemed appropriate due to the specified multiple hurdle risk reduction approach outlined. Any non synthetic crop treatment that contains animal manure must use only fully composted manure in addition to a validated process and pass testing requirements before a application to soils or directly to edible portions of lettuce and leafy greens.</li> </ul>

566

567  
568  
569  
570

**Figure 3. Decision Tree for Nonsynthetic Crop Treatments That Contain Animal Products**



571  
572

573 **Note: Mixtures of soil amendment materials**

574

575 For soil amendments that contain mixtures of materials each component must meet the  
576 requirements of its respective class of materials. The usages allowed will conform to that of the  
577 most stringent class of materials utilized in the mixture.

578

579 For example; Soil amendments containing animal manure that has been physically heat treated  
580 or processed by other equivalent methods mixed with soil amendments not containing animal  
581 manure would require a process certification for the physically heat treated or processed by  
582 other equivalent methods materials and the components from non-animal manure would  
583 require documentation attesting to its manure free status. The resulting mixture could then be  
584 applied in accordance with the guidelines associated with the physically heated treated class of  
585 materials (most stringent limits).

586 **8. ISSUE: HARVEST EQUIPMENT**

587 This section addresses harvest and harvest aid equipment used for lettuce/leafy greens.  
588 Mechanical or machine harvest has become increasingly prevalent and provides opportunity  
589 for increased surface contact exposure. This includes field cored lettuce operations that use  
590 various harvest equipment and aids.

591

592 **8.1. The Best Practices Are:**

- 593 • Prepare an SOP for harvest equipment that addresses the following:
- 594 ○ Sanitation verification
- 595 ○ Daily inspection
- 596 ○ Periodic microbial swabs or other equivalent indicator
- 597 ○ Proper cleaning, sanitation and storage of hand harvest equipment (knives,  
598 scythes, etc.)
- 599 ○ Control procedures when equipment is not in use, including policy for  
600 removal of equipment from the work area or site and the use of scabbards,  
601 sheathes or other storage equipment.
- 602 • Prepare an SOP for handling and storage of product containers that addresses the  
603 following:
- 604 ○ Over night storage
- 605 ○ Contact with the ground
- 606 ○ Container assembly (RPC, fiber bin, plastic bin, etc)
- 607 ○ Damaged containers
- 608 ○ Use of containers only as intended
- 609 • Prepare an SOP for sanitary operation of equipment which addresses:.
- 610 ○ Spills and leaks
- 611 ○ Inoperative water sprays

- 612                   ○ Exclusion of foreign objects (including glass, plastic, metal and other
- 613                   debris)
- 614                   ○ Establish and implement cleaning and sanitation schedules for containers
- 615                   and equipment that will be used in hydration.
- 616                   ○ Maintain logs documenting cleaning and sanitation, and retain these
- 617                   records for at least two years.
- 618                   ○ Establish policies for the storage and control of water tanks and
- 619                   equipment used for hydration operations when not in use.
- 620
- 621                   ● Establish appropriate measures that reduce and control the potential introduction
- 622                   of human pathogens at the cut surface during and after mechanical harvest
- 623                   operations. Due to the cut surface being more vulnerable to microbial
- 624                   contamination, this best practice is extremely important and all practical means
- 625                   should be taken to reduce the possibility of introduction of contamination at this
- 626                   process step.
- 627                   ● If re-circulated rinse or antioxidant solutions are used on the cut surface, take all
- 628                   practicable precautions to prevent them from becoming a source of
- 629                   contamination.
- 630                   ● Design equipment to facilitate cleaning by using materials and construction that
- 631                   facilitate cleaning and sanitation of equipment food contact surfaces (e.g.,
- 632                   transportation tarps, conveyor belts, etc.).
- 633                   ● Establish the frequency of equipment cleaning and sanitation by developing
- 634                   Sanitation Standard Operating Procedures (SSOPs) and a sanitation schedule for
- 635                   machine harvest operations.
- 636                   ● Evaluate the use of cleaning verification methods for harvesting equipment (e.g.,
- 637                   ATP test methods).
- 638                   ● Locate equipment cleaning and sanitizing operations away from product and other
- 639                   equipment to reduce the potential for cross contamination.
- 640                   ● Establish equipment storage and control procedures to minimize the potential for
- 641                   contamination when not in use. Establish policies and sanitary design options that
- 642                   facilitate frequent and thorough cleaning and sanitizing of food contact surfaces.
- 643                   ● Develop and implement appropriate cleaning, sanitizing, storage and handling
- 644                   procedures of all food contact surfaces to reduce and control the potential for
- 645                   microbial cross contamination.
- 646                   ● Allow adequate distance for the turning and manipulation of harvest equipment to
- 647                   prevent cross contamination from areas of animal or significant risk intrusion or
- 648                   adjacent land that may pose a risk.
- 649

650 **9. ISSUE: HARVEST PERSONNEL - DIRECT CONTACT WITH SOIL DURING HARVEST**  
651 **(FIELD SANITATION)**

652 After manual harvest of lettuce/leafy greens, placing or stacking product on soil before the  
653 product is placed into a container may expose the product to human pathogens if the soil is  
654 contaminated. Research has demonstrated that microbes, including human pathogens, can  
655 readily attach to cut lettuce/leafy green surfaces (Takeuchi *et al.* 2001).  
656

657 **9.1. The Best Practices Are:**

- 658 • Evaluate appropriate measures that reduce and control the potential introduction  
659 of human pathogens through soil contact at the cut surface after harvest (e.g.  
660 frequency of knife sanitation, no placement of cut surfaces of harvested product  
661 on the soil, container sanitation, single use container lining, etc.).
- 662 • Do not stack soiled bins on top of each other if the bottom of one bin has had  
663 direct contact with soil unless a protective barrier (*i.e.*, liner, cover, *etc.*) is used  
664 to separate the containers..  
665

666 **10. ISSUE: FIELD AND HARVEST PERSONNEL - TRANSFER OF HUMAN PATHOGENS**  
667 **BY WORKERS (FIELD SANITATION)**

668 Lettuce/leafy greens are handled by harvest crews during harvest in that each lettuce/leafy  
669 greens plant is touched/handled as part of the harvest process. It is possible that persons  
670 working with produce in the field may transfer microorganisms of significant public health  
671 concern. Workers may be asymptomatic.

672 **10.1. The Best Practices Are:**

- 673 • Use appropriate preventive measures outlined in GAPs such as training in appropriate  
674 and effective hand washing, glove use and replacement, and mandatory use of  
675 sanitary field latrines to reduce and control potential contamination.
- 676 • Establish a written worker practices program (*i.e.*, an SOP) that can be used to verify  
677 employee compliance with company food safety policy. This program shall establish  
678 the following practices for field and harvest employees as well as visitors.
  - 679 ○ Prior to harvest, an individual should be designated as responsible for  
680 harvesting food safety
  - 681 ○ Use, storage, record keeping, and proper labeling of chemicals
  - 682 ○ Training on proper sanitation and hygiene practices
  - 683 ○ Requirements for workers to wash their hands before beginning or returning  
684 to work
  - 685 ○ Confinement of smoking, eating and drinking of beverages other than water  
686 to designated areas.
  - 687 ○ Personal item storage

- 688 • A written physical hazard prevention program should be developed for leafy green  
689 products that are intended for further processing. The program must address the  
690 following:
  - 691 ○ Employee clothing and jewelry (head and hair restraints, aprons, gloves,  
692 visible jewelry, etc.)
  - 693 ○ Removal of all objects from upper pockets
  - 694 ○ Foreign objects in the field.
- 695 • Establish a worker health practices program (i.e., an SOP) that address the following  
696 issues:
  - 697 ○ Workers with diarrhea disease or symptoms of other infectious disease are  
698 prohibited from handling fresh produce.
  - 699 ○ Workers with open cuts or lesions are prohibited from handling fresh produce  
700 without specific measures to prevent cross contamination of product.
  - 701 ○ Actions for employee to take in the event of injury or illness.
  - 702 ○ A policy describing procedures for handling/disposition of produce or food  
703 contact surfaces that have come into contact with blood or other body fluids.
- 704 • A field sanitary facility program (i.e., an SOP) shall be implemented, and it should  
705 address the following issues: the number, condition, and placement of field sanitation  
706 units, the accessibility of the units to the work area, facility maintenance, facility  
707 supplies (i.e., hand soap, water, paper towels, toilet paper, etc.), facility signage,  
708 facility cleaning and servicing, and a response plan for major leaks or spills.
  - 709 ○ Sanitary facilities should be placed such that the location minimizes the  
710 impact from potential leaks and/or spills while allowing access for cleaning  
711 and service.
  - 712 ○ The location and sanitary design of toilets and hand wash facilities should be  
713 optimized to facilitate the control, reduction and elimination of human  
714 pathogens from employee hands. Evaluate the location of worker hygiene  
715 facilities to maximize accessibility and use, while minimizing the potential  
716 for the facility to serve as a source of contamination.
  - 717 ○ Establish the frequency of toilet and hand washing facility  
718 maintenance/sanitation.
  - 719 ○ Establish equipment and supply storage and control procedures when not in  
720 use.
  - 721 ○ Maintain documentation of maintenance and sanitation schedules and any  
722 remedial practices for a period of two years.

723 **11. ISSUE: EQUIPMENT FACILITATED CROSS CONTAMINATION (FIELD**  
724 **SANITATION)**

725 When farm equipment has had direct contact with raw untreated manure, untreated compost,  
726 waters of unknown quality, animals of significant risk, or other potential human pathogen  
727 reservoirs it may be a source of cross contamination. Such equipment should not be used in  
728 proximity to or in areas where it may contact edible portions of lettuce and or leafy greens.

729

730 **11.1. The Best Practices Are:**

- 731 • Identify any field operations that may pose a risk for cross-contamination. These  
732 include management personnel in the fields, vehicles used to transport workers,  
733 as well as many other possibilities.
- 734 • Segregate equipment used in high-risk operations or potentially exposed to high  
735 levels of contamination.
- 736 • Use effective means of equipment cleaning and sanitation before subsequent  
737 equipment use in lettuce/leafy greens production, if it was previously used in a  
738 high-risk operation.
- 739 • Develop appropriate means of reducing and controlling the possible transfer of  
740 human pathogens to soil and water that may directly contact edible lettuce/leafy  
741 green tissues through use of equipment.
- 742 • Maintain appropriate records related to equipment cleaning and possible cross-  
743 contamination issues for a period of two years.

744

745 **12. ISSUE: FLOODING**

746 Flooding for purposes of this document is defined as the flowing or overflowing of a field  
747 with water outside of a producer's control, that is reasonably likely to contain  
748 microorganisms of significant public health concern and is reasonably likely to cause  
749 adulteration of the edible portions of fresh produce in that field. Pooled water (e.g., rainfall)  
750 that is not reasonably likely to contain microorganisms of significant public health concern  
751 and is not reasonably likely to cause adulteration of the edible portion of fresh produce  
752 should not be considered flooding.

753

754 If flood waters contain microorganisms of significant public health concern, crops in close  
755 proximity to soil such as lettuce/leafy greens may be contaminated if there is direct contact  
756 between flood water or contaminated soil and the edible portions of lettuce/leafy greens  
757 (Wachtel *et al.* 2002a;2002b).

758

759 In the November 4, 2005 FDA "Letter to California Firms that Grow, Pack, Process, or Ship  
760 Fresh and Fresh-cut Lettuce/leafy greens" the agency stated that it "considers ready to eat  
761 crops (such as lettuce/leafy greens) that have been in contact with flood waters to be  
762 adulterated due to potential exposure to sewage, animal waste, heavy metals, pathogenic  
763 microorganisms, or other contaminants. FDA is not aware of any method of reconditioning  
764 these crops that will provide a reasonable assurance of safety for human food use or  
765 otherwise bring them into compliance with the law. Therefore, FDA recommends that such  
766 crops be excluded from the human food supply and disposed of in a manner that ensures they  
767 do not contaminate unaffected crops during harvesting, storage or distribution.

768

769 "Adulterated food may be subject to seizure under the Federal Food, Drug, and Cosmetic  
770 Act, and those responsible for its introduction or delivery for introduction into interstate  
771 commerce may be enjoined from continuing to do so or prosecuted for having done so. Food

772 produced under unsanitary conditions whereby it may be rendered injurious to health is  
773 adulterated under § 402(a)(4) of the Federal Food, Drug, and Cosmetic Act (21 U.S.C. 342(a)  
774 (4); (US FDA 2004).

775  
776 Areas that have been flooded can be separated into three groups: 1) product that has come  
777 into contact with flood water, 2) product that is in proximity to a flooded field but has not  
778 been contacted by flood water, and 3) production ground that was partially or completely  
779 flooded in the past before a crop was planted. The considerations for each situation are  
780 described below and presented in Table 4.

781

782 **12.1. The Best Practices For Product That Has Come Into Contact With**  
783 **Flood Water Are:**

- 784 • See Table 4 for numerical criteria for lettuce and leafy greens production fields  
785 that have possibly come into contact with flood waters. The “Technical Basis  
786 Document” (Appendix B) describes the process used to develop these metrics.
- 787 • FDA considers any crop that has come into contact with floodwater to be an  
788 “adulterated” commodity that cannot be sold for human consumption.
- 789 • To reduce the potential for cross contamination do not drive harvest equipment  
790 through flooded areas reasonably likely to contain microorganisms of public  
791 health significance (see previous section).

792

793  
794

**TABLE 4. FLOODING**

When evidence of flooding in a production block occurs.

Practice	Metric/Rationale
<b>Flooding Defined</b>	The flowing or overflowing of a field with water outside a producer’s control that is reasonably likely to contain microorganisms of significant public health concern and is reasonably likely to cause adulteration of edible portions of fresh produce in that field. Additional discussion of this definition and implications for production is provided in the text portion of this document.
<b>Allowable Harvest Distance from Flooding</b>	<ul style="list-style-type: none"> <li>• Buffer and do not harvest any product within 30 ft of the flooding.</li> <li>• Required buffer distance may be greater than 30 ft based on risk analysis by food safety professional.</li> <li>• If there is evidence of flooding, the production block must undergo a detailed food safety assessment by appropriately trained food safety personnel (see Glossary) prior to harvest, as defined in the text of this document.</li> </ul>
<b>Verification</b>	<ul style="list-style-type: none"> <li>• Documentation must be archived for a period of two years following the flooding event. Documentation may include photographs, sketched maps, or other means of delineating affected portions of production fields.</li> </ul>
<b>Time Interval Before Planting Can Commence Following the Receding of Floodwaters</b>	<ul style="list-style-type: none"> <li>• 60 days prior to planting provided that the soil has sufficient time to dry out.</li> <li>• Appropriate soil testing can be used to shorten this period to 30 days prior to planting. This testing must be performed in a manner that accurately represents the production field and indicates soil levels of microorganisms lower than the recommended standards for processed compost. Suitable representative samples should be collected for the entire area suspected to have been exposed to flooding. For additional guidance on appropriate soil sampling techniques, use the <i>Soil Screening Guidance: Technical Background Document</i> (US EPA 1996). Specifically, Part 4 provides guidance for site investigations. Reputable third-party environmental consultants or laboratories provide sampling services consistent with this guidance.</li> <li>• Appropriate mitigation and mitigation strategies are included in the text portion of the document.</li> </ul>
<b>Rationale</b>	<ul style="list-style-type: none"> <li>• <a href="#">The basis for the 30 foot distance is the turn around distance for production equipment to prevent cross-contamination of non-flooded ground or produce.</a></li> </ul>

795  
796

797 **12.2. The Best Practices for Product in Proximity to a Flooded Area But**  
798 **Not Contacted By Flood Water Are:**

- 799 • Prevent cross contamination between flooded and non-flooded areas (e.g.  
800 cleaning equipment, eliminating contact of any farming or harvesting equipment  
801 or personnel with the flooded area during growth and harvest of non-flooded  
802 areas).
- 803 • To facilitate avoiding contaminated/adulterated produce, place markers  
804 identifying both the high-water line of the flooding and an interval 30 feet beyond  
805 this line. If 30 feet is not sufficient to prevent cross contamination while turning  
806 harvesting or other farm equipment in the field, use a greater appropriate interval.  
807 Take photographs of the area for documentation. Do not harvest product within  
808 the 30 foot buffer zone.

809

810 **12.3. The Best Practices For Formerly Flooded Production Ground Are:**

- 811 • Allow soils to dry sufficiently and be reworked prior to planting subsequent crops  
812 on formerly flooded production ground.
- 813 • Do not replant formerly flooded production ground for at least 60 days following  
814 the receding of floodwaters. This period or longer and active tillage of the soil  
815 provide additional protection against the survival of pathogenic organisms.
- 816 • If flooding has occurred in the past on the property, soil clearance testing may be  
817 conducted prior to planting leafy greens. Soil testing may be used to shorten the  
818 clearance period to 30 days. If performed, testing must indicate soil levels of  
819 microorganisms lower than the standards for processed compost. Suitable  
820 representative samples should be collected for the entire area suspected to have  
821 been exposed to flooding.
- 822 • Sample previously flooded soil for the presence of microorganisms of significant  
823 public health concern or appropriate indicator microorganisms. Microbial soil  
824 sampling can provide valuable information regarding relative risks; however,  
825 sampling by itself does not guarantee that crops grown within the formerly  
826 flooded production area will be free of the presence of human pathogens.
- 827 • Prior to replanting or soil testing, the designated food safety professional for the  
828 producer shall perform a detailed food safety assessment of the production field.  
829 This designated professional will be responsible for assessing the relative merits  
830 of testing versus observing the appropriate time interval for planting, and also  
831 will coordinate any soil testing plan with appropriate third-party consultants  
832 and/or laboratories that have experience in this type of testing.
- 833 • Evaluate the field history and crop selection on formerly flooded production  
834 ground.
- 835 • Assess the time interval between the flooding event, crop planting, and crop  
836 harvest. Comparative soil samples may be utilized to assess relative risk if

837 significant reductions in indicator microorganisms have occurred within this time  
838 interval.

839 • Evaluate the source of flood waters (e.g., drainage canal, river, irrigation canal,  
840 etc.) for potential significant upstream contributors of human pathogens at levels  
841 that pose a significant threat to human health.

842 • Prevent cross-contamination by cleaning or sanitizing any equipment that may  
843 have contacted previously flooded soil (also see the section on Equipment  
844 Facilitated Cross Contamination above).

845 **13. ISSUE: PRODUCTION LOCATIONS - CLIMATIC CONDITIONS AND ENVIRONMENT**

846 Lettuce/leafy greens are grown in varying regions but generally in moderate weather  
847 conditions. Cool, humid conditions favor human pathogen persistence (Takeuchi and Frank  
848 2000; Takeuchi *et al.* 2000) while drier climates may present other problems such as  
849 requirements for additional water that may increase the potential for introduction of human  
850 pathogens. Heavy rains in certain areas may also cause lettuce/leafy greens to be exposed to  
851 contaminated soil due to rain splashing. It is important to tailor practices and procedures  
852 designed to promote food safety to the unique environment in which each crop may be  
853 produced  
854

855 **13.1. The Best Practices Are:**

856 • Consider harvest practices such as removing soiled leaves, not harvesting soiled  
857 heads, etc., when excessive soil or mud builds up on lettuce/leafy greens.

858 • Take care to reduce the potential for windborne soil, including soil from roads  
859 adjacent to fields, water, or other media that may be a source of contamination to  
860 come into direct contact with the edible portions of lettuce and leafy greens. Do not  
861 allow runoff from adjacent properties to come into contact with produce.

862 • Evaluate and implement practices to reduce the potential for the introduction of  
863 pathogens into production blocks by wind or runoff. Such practices may include but  
864 are not limited to berms, windbreaks, diversions ditches and vegetated filter strips.

865 • When soil has accumulated on plants, remove soil during the harvest or further  
866 processing.

867

868 **14. ISSUE: PRODUCTION LOCATIONS - ENCROACHMENT BY ANIMALS AND URBAN**  
869 **SETTINGS**

870 Lettuce/leafy greens are generally grown in rural areas that may have adjacent wetlands,  
871 wildlands, and/or parks harboring wildlife. Some wildlife species are known to be potential  
872 carriers of various human pathogens (Fenlon 1985). Specific wildlife species that have been  
873 shown to pose the greatest risk are the focus of this section and are listed in Table 5. In  
874 addition, extensive development in certain farming communities has also created situations  
875 with urban encroachment and unintentional access by domestic animals and livestock which  
876 may also pose varying degrees of risk depending on the animal species. Finally, it is possible  
877 that some land uses may be of greater concern than others when located near production  
878 fields. Table 6 provides a list of these uses and recommended buffer distances.

879

880 **14.1. The Best Practices Are:**

- 881 • See Tables 5 and 6 and Decision Tree (Figure 5) for numerical criteria and  
882 guidance applicable to animal encroachment and adjacent land uses. The  
883 “Technical Basis Document” (Appendix B) describes the process used to develop  
884 these metrics.
- 885 • During the Environmental Assessments discussed in Section 2, the location of  
886 any adjacent land uses that may be of potential risk should be documented. In  
887 addition, as specified in Table 6, any deviations from the recommended buffer  
888 distances due to mitigation factors or increased risk should be documented and  
889 explained.
- 890 • Fencing, vegetation removal, and destruction of habitat may result in adverse  
891 impacts to the environment. Potential adverse impacts include loss of habitat to  
892 beneficial insects and pollinators; wildlife loss; increased discharges of sediment  
893 and other pollutants resulting from the loss of vegetative filtering; and increased  
894 air quality impacts if bare soil is exposed to wind. It is recommended that  
895 producers check for local, state, and federal laws and regulations that protect  
896 riparian habitat, restrict removal of vegetation or habitat, or restrict construction  
897 of wildlife deterrent fences in riparian areas or wildlife corridors.
- 898 • Document any observed encroachment by animals of significant risk during  
899 production periods.
- 900 • Evaluate and monitor animal of significant risk activity in and proximate to  
901 lettuce/leafy greens fields and production environments. Conduct periodic  
902 monitoring, pre-season, pre-harvest, and harvest assessments. If there are  
903 animals of significant risk present, make particular efforts to reduce their access  
904 to lettuce and leafy green produce.
- 905 • Evaluate the risk to subsequent crop production on production acreage that has  
906 experienced recent postharvest grazing with or by domesticated animals that used  
907 field culls as a source of animal feed.
- 908 • Locate production blocks to minimize potential access by animals of significant  
909 risk and maximize distances to possible sources of microbial contamination. For  
910 example, consider the proximity to water (i.e., riparian areas), animal of  
911 significant risk harborage, open range lands, non-contiguous blocks, urban  
912 centers, etc. Periodically monitor these factors and assess during pre-season and  
913 preharvest assessments as outlined in Tables 5 and 6. If the designated food  
914 safety professional deems that there is the potential for microbial contamination  
915 from adjacent areas, a risk assessment shall be performed to determine the risk  
916 level as well as to evaluate potential strategies to control or reduce the  
917 introduction of human pathogens.
- 918 • DO NOT harvest areas of fields where unusually heavy activity by animals of  
919 significant risk occurs. If animal of significant risk intrusions are common on a  
920 particular production field, consider fencing, barriers, noisemakers, and other  
921 practices that may reduce intrusions.

- 922 • Train harvest employees to recognize and report evidence (e.g., feces) of animal  
923 of significant risk activity.
- 924 • Pooled water (e.g., a seasonal lake) from rainfall may attract animals of  
925 significant risk and should be considered as part of any land use evaluation.
- 926 • Consider controlling risks associated with encroachment by urban development.  
927 Risks may include, but are not limited to, domestic animal fecal contamination of  
928 production fields and harvest equipment and septic tank leaching.
- 929 • Producers are encouraged to contact the relevant agencies (e.g., the Regional  
930 Water Quality Control Board and state and federal fish and wildlife agencies) to  
931 confirm the details of these requirements. In addition, producers may wish to  
932 consult with local NRCS to evaluate the food safety risks associated with wildlife  
933 of significant risk, livestock, domestic animals and other adjacent land uses and  
934 to develop and document strategies to control or reduce the introduction of  
935 human pathogens through wildlife of significant risk for each production block.

936  
937

**TABLE 5. ANIMAL OF SIGNIFICANT RISK ACTIVITY IN FIELD (WILD OR DOMESTIC)**

When evidence of animal of significant risk intrusion in a production block occurs.

Issue	Metric	Remedial Actions
<p><b>Evidence of Intrusion</b></p>	<p><u>Frequency</u></p> <ul style="list-style-type: none"> <li>• There shall be a periodic monitoring plan in place for production fields.</li> <li>• There shall be Pre Season, Pre Harvest, and Harvest Assessments</li> </ul> <p><u>Variables</u></p> <ul style="list-style-type: none"> <li>• Physical observation of animals in the field</li> <li>• Downed fences</li> <li>• Animal tracks in production block</li> <li>• Animal feces or urine in production block</li> <li>• Eaten plants in production block</li> </ul> <p><u>Animals of Significant Risk</u></p> <ul style="list-style-type: none"> <li>• Deer</li> <li>• Pigs (wild and domestic)</li> <li>• Cattle</li> <li>• Goats and Sheep</li> </ul>	<ul style="list-style-type: none"> <li>• If there is evidence of intrusion by animals of significant risk, the production block must undergo a detailed food safety assessment by appropriately trained food safety personnel (see Glossary) prior to harvest, as defined in the text of this document.</li> <li>• In developing remedial and corrective actions, consider consulting with wildlife and/or domestic animal experts as appropriate.</li> <li>• If remedial actions cannot be formulated that control or eliminate the identified risk, destroy the block by disking under the crop.</li> <li>• Equipment used to destroy crop must be cleaned and sanitized upon exiting the field.</li> <li>• Investigate potential causes for intrusion by animals of significant risk and assess the extent of intrusion and impact on crop food risk.</li> <li>• Formulate effective corrective actions. Prior to taking action that may affect natural resources, producers should check local, state and federal laws and regulations that protect riparian habitat, restrict removal of vegetation or habitat, or restrict construction of wildlife deterrent fences in riparian areas or wildlife corridors.</li> <li>• Evidence of intrusion by animals of significant risk and corrective actions shall be documented and available for verification for a period of two years.</li> </ul>
<p><b>Allowable Harvest Distance from Evidence of Intrusion</b></p>	<p><b>Please see Figure 5. Decision Tree for Conducting Pre-Harvest and Harvest Assessments.</b></p> <p><u>Monitoring</u> Evaluate and monitor animal of significant risk activity in and proximate to lettuce/leafy greens fields and production environments. Conduct periodic monitoring, pre-season, pre-harvest, and harvest assessments.</p> <p><u>Pre Harvest Assessment:</u> Conduct the pre-harvest assessment not more than one week prior to harvest.</p> <p><b>Fecal Material</b></p> <ul style="list-style-type: none"> <li>• Do not harvest any produce that has come into direct contact with fecal material.</li> <li>• If evidence of fecal material is found, conduct a food safety assessment using qualified personnel. Do not harvest any crop found</li> </ul>	

Issue	Metric	Remedial Actions
	<p>within a minimum 5 foot radius buffer distance from the spot of the contamination unless remedial action can be found that adequately control the risk. The food safety professional can increase this buffer distance if deemed appropriate.</p> <ul style="list-style-type: none"> <li>Remove fecal material from the field and dispose of properly.</li> </ul> <p><b>Intrusion</b></p> <ul style="list-style-type: none"> <li>If evidence of animal of significant risk intrusion is found in a production field, conduct a visual food safety assessment to determine whether the areas of intrusion can be adequately controlled (e.g., solitary deer track with no evidence of feeding), or whether a three foot buffer radius non-harvest area should be applied (e.g., wide areas of wild pig rooting and tracks).</li> </ul> <p><i>Harvest Assessment</i></p> <p>If evidence of animal of significant risk intrusion into the production block is not discovered until harvest operations:</p> <ul style="list-style-type: none"> <li>Stop harvest operations.</li> <li>Initiate an intensified block assessment for evidence of further contamination and take appropriate actions per the aforementioned actions.</li> <li>If evidence of intrusion is discovered during production block harvest operations and the harvest rig has been potentially contaminated by contaminated product or feces, clean and sanitize the equipment before resuming harvest operations.</li> <li>Require all employees to wash and sanitize their hands/gloves before resuming harvest operations.</li> <li>If contamination is discovered in harvest containers such as bins/totes, discard the product, and clean and sanitize the container before reuse.</li> </ul>	
<b>Verification</b>	<ul style="list-style-type: none"> <li>Archive documentation for a period of two years following the intrusion event. Documentation may include photographs, sketched maps, or other means of delineating affected portions of production fields.</li> </ul>	
<b>Rationale</b>	<ul style="list-style-type: none"> <li>The basis of these metrics is qualitative assessment of the relative risk from a variety of intrusions. Some animal feces and some signs of intrusion (feces vs. tracks) are considered to be of more concern than others. Because it is difficult to develop quantitative metrics for these types of risks, a food safety assessment is considered appropriate for this issue.</li> </ul>	

938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948

**TABLE 6. CROP LAND AND WATER SOURCE ADJACENT LAND USE**


Land Use/Water Source	Metric (This distance may be either increased or decreased depending on risk and mitigation factors.)	Considerations for Risk Analysis*		
		Risk/Mitigation Factors	Increase Distance	Decrease Distance
Composting Operations (manure or animal products)	Due to the lack of science at this time, an interim guidance distance of 400 ft from the edge of crop is proposed. This number is subject to change as science becomes available.  The proximate safe distance depends on the risk/mitigation factors listed to the right. Evaluate risk and document consideration of these factors. Research is being proposed to study appropriate distance.	Distance from active compost operation	--	--
		Topography: Uphill from crop	√	
		Topography: Downhill from crop		√
		Opportunity for water run off through or from composting operations	√	
		Opportunity for soil leaching	√	
		Presence of physical barriers such as windbreaks, diversion ditches, vegetative strips		√
Concentrated Animal Feeding Operations (as defined in 40 CFR 122.23)	Due to the lack of science at this time, an interim guidance distance of 400 ft from the edge of crop is proposed. This number is subject to change as science becomes available.  The proximate safe distance depends on the risk/mitigation factors listed to the right. Evaluate risk and document consideration of these factors. Research is being proposed to study appropriate distance.	Fencing and other physical barriers such as berms, diversion ditches and vegetated strips can be employed to prevent intrusion of domestic animals, control runoff, etc.		√
		Topography: Uphill from crop	√	
		Topography: Downhill from crop		√
		Opportunity for water run off through or from CAFOs	√	
		Opportunity for soil leaching	√	
		Manure Management Program utilized		√
Non-synthetic Soil Amendment Pile (containing manure or animal products)	Due to the lack of science at this time, an interim guidance distance of 400 ft from the edge of crop is proposed. This number is subject to change as science becomes available.  The proximate safe distance depends on the risk/mitigation factors listed to the right. Evaluate risk and document consideration of these factors. Research is being proposed to study appropriate distance.	Access and review COA for materials in question.		√
		Topography: Uphill from crop	√	
		Topography: Downhill from crop		√
		Opportunity for water run off through or from non-synthetic soil amendment storage areas	√	
		Opportunity for soil leaching	√	

Land Use/Water Source	Metric (This distance may be either increased or decreased depending on risk and mitigation factors.)	Considerations for Risk Analysis*		
		Risk/Mitigation Factors	Increase Distance	Decrease Distance
	For non-synthetic crop treatments that have been heat treated using a validated process an interim guidance distance of 30 feet from the edge of the crop is proposed	Covering on pile to prevent wind dispersion		√
Grazing Lands/Domestic Animals (includes homes with hobby farms, and non commercial livestock)	30 ft from the edge of crop.	Fencing and other physical barriers such as berms, diversion ditches and vegetated strips can be employed to prevent intrusion of domestic animals, control runoff, etc.		√
		Topography: Uphill from crop	√	
		Topography: Downhill from crop		√
		Opportunity for water run off through or from grazing lands	√	
		Opportunity for soil leaching	√	
Homes or other building with a septic leach field.	30 ft from the edge of crop to the leach field.	Active leach field: < 10 yrs old		√
		Active leach field: > 25 yrs old	√	
		Inactive leach field		√
		Topography: Uphill from crop	√	
		Topography: Downhill from crop		√
		Physical barriers		√
Well Head Distance from Untreated Manure	200 ft separation of untreated manure from wells, although less distance may be sufficient.	Topography: Uphill from manure		√
		Topography: Downhill from manure	√	
		Opportunity for water run off from or through untreated manure to well head	√	
		Opportunity for soil leaching	√	
		Presence of physical barriers such as windbreaks, diversion ditches, vegetative strips		√
Surface Water Distance from	At least 100 feet separation for sandy soil and 200 feet	Topography: Uphill from manure		√

Land Use/Water Source	Metric (This distance may be either increased or decreased depending on risk and mitigation factors.)	Considerations for Risk Analysis*		
		Risk/Mitigation Factors	Increase Distance	Decrease Distance
Untreated Manure	separation for loamy or clay soil (slope less than 6%; increase distance to 300 feet if slope greater than 6%) is recommended.	Topography: Downhill from manure	√	
		Opportunity for water runoff from or through untreated manure to surface waters.	√	
		Opportunity for soil leaching	√	
		Presence of physical barriers such as windbreaks, diversion ditches, vegetative strips		√
Rationale	<ul style="list-style-type: none"> <li>The bases for these distances above is best professional judgment of authors, contributors, and expert reviewers to prevent potential cross-contamination from adjacent land uses, taking into consideration the 200 foot distance cited in FDA (US FDA 2001) for separation of manure from wellheads and the 30 foot turn-around distance for production equipment. Because of the numerous factors that must be taken into account to determine appropriate distances, a qualitative assessment of the relative risk from various types of land use and surface waters was used to determine appropriate distances.</li> </ul>			

951 Producers should check for local, state and federal laws and regulations that protect riparian habitat, restrict removal of vegetation or habitat, or restrict construction of  
952 wildlife deterrent fences in riparian areas or wildlife corridors. Producers may want to contact the relevant agencies (e.g., the Regional Water Quality Control Board  
953 and state and federal fish and wildlife agencies) to confirm the details of these requirements.

954 **Figure 5. Decision Tree for Conducting Pre-harvest and Harvest Assessment of Animal Activity in Field**  
 955 **(Wild or Domestic)**  
 956



957  
 958  
 959  
 960

961 15. DETAILED BACKGROUND GUIDANCE INFORMATION

962

963 15.1. **Required Reference Documents**

964

- 965 1. FDA Guide to Minimize Microbial Food Safety Hazards for Fresh Fruits and Vegetables  
966 (www.foodsafety.gov/~dms/prodguid.html)
- 967 2. UFFVA Food Safety Auditing Guidelines: Core Elements of Good Agricultural Practices for Fresh  
968 Fruits and Vegetables
- 969 3. UFFVA Food Safety Questionnaire for Fresh Fruits and Vegetables
- 970 4. National GAPs Program Cornell University: Food Safety Begins on the Farm: A Grower Self  
971 Assessment of Food Safety Risks  
972

973 15.2. **References**

974

- 975 CCR Title 14 - Chapter 3.1 - Article 5. 2007. *Article 5. Composting Operation and Facility Siting and*  
976 *Design Standards*. Accessed February 15, 2007.  
977 <http://www.ciwmb.ca.gov/regulations/Title14/ch31a5.htm#article5>
- 978 Fukushima H, Hoshina K, and Gomyoda M. 1999. Long-term survival of shiga toxin-producing  
979 Escherichia coli O26, O111, and O157 in bovine feces. *Applied and environmental microbiology*  
980 65 (11):5177-81.  
981 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=10543842)  
982 [uids=10543842](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=10543842)
- 983 Gagliardi JV and Karns JS. 2000. Leaching of Escherichia coli O157:H7 in diverse soils under various  
984 agricultural management practices. *Applied and environmental microbiology* 66 (3):877-83.  
985 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=10698745)  
986 [uids=10698745](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=10698745)
- 987 Islam M, Doyle MP, Phatak SC, Millner P, and Jiang X. 2004. Persistence of enterohemorrhagic  
988 Escherichia coli O157:H7 in soil and on leaf lettuce and parsley grown in fields treated with  
989 contaminated manure composts or irrigation water. *Journal of food protection* 67 (7):1365-70.  
990 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=15270487)  
991 [uids=15270487](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=15270487)
- 992 Jiang X, Morgan J, and Doyle MP. 2003. Fate of Escherichia coli O157:H7 during composting of bovine  
993 manure in a laboratory-scale bioreactor. *Journal of food protection* 66 (1):25-30.  
994 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=12540177)  
995 [uids=12540177](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=12540177)
- 996 Solomon EB, Pang HJ, and Matthews KR. 2003. Persistence of Escherichia coli O157:H7 on lettuce  
997 plants following spray irrigation with contaminated water. *Journal of food protection* 66  
998 (12):2198-202.  
999 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=14672213)  
1000 [uids=14672213](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=14672213)
- 1001 Stine SW, Song I, Choi CY, and Gerba CP. 2005. Application of microbial risk assessment to the  
1002 development of standards for enteric pathogens in water used to irrigate fresh produce. *Journal of*  
1003 *food protection* 68 (5):913-8.  
1004 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=15895721)  
1005 [uids=15895721](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=15895721)

- 1006 Suslow, T.V., M.P. Oria, L.R. Beuchat, E.H. Garrett, M.E. Parish, L.J. Harris, J.N. Farber, F.F. Busta.  
1007 2003. Production practices as risk factors in microbial food safety of fresh and fresh-cut produce.  
1008 *Comprehensive Reviews in Food Science and Food Safety* 2S:38-77.
- 1009 Takeuchi K and Frank JF. 2000. Penetration of Escherichia coli O157:H7 into lettuce tissues as affected  
1010 by inoculum size and temperature and the effect of chlorine treatment on cell viability. *Journal of*  
1011 *food protection* 63 (4):434-40.  
1012 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10772206)  
1013 [uids=10772206](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10772206)
- 1014 Takeuchi K, Matute CM, Hassan AN, and Frank JF. 2000. Comparison of the attachment of Escherichia  
1015 coli O157:H7, Listeria monocytogenes, Salmonella typhimurium, and Pseudomonas fluorescens to  
1016 lettuce leaves. *Journal of food protection* 63 (10):1433-7.  
1017 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11041147)  
1018 [uids=11041147](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11041147)
- 1019 Takeuchi K, Hassan AN, and Frank JF. 2001. Penetration of Escherichia coli O157:H7 into lettuce as  
1020 influenced by modified atmosphere and temperature. *Journal of food protection* 64 (11):1820-3.  
1021 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11726166)  
1022 [uids=11726166](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11726166)
- 1023 US EPA. 1996. *Soil Screening Guidance: Technical Background Document*. EPA/540/R95/128: Office of  
1024 Solid Waste and Emergency Response, United States Environmental Protection Agency.  
1025 [http://rais.ornl.gov/homepage/SSG\\_nonrad\\_technical.pdf](http://rais.ornl.gov/homepage/SSG_nonrad_technical.pdf)
- 1026 US EPA. 2002. *Implementation Guidance for Ambient Water Quality Criteria for Bacteria: May 2002*  
1027 *Draft*. EPA-823-B-02-003: United States Environmental Protection Agency.  
1028 <http://www.epa.gov/waterscience/standards/bacteria/bacteria.pdf>
- 1029 US FDA. 2001. Chapter II: Production Practices as Risk Factors in Microbial Food Safety of Fresh and  
1030 Fresh-Cut Produce. In *Analysis and Evaluation of Preventive Control Measures for the Control*  
1031 *and Reduction/Elimination of Microbial Hazards on Fresh and Fresh-Cut Produce*; pp.  
1032 <http://www.cfsan.fda.gov/~comm/ift3-2a.html>.
- 1033 US FDA. 2004. Federal Food, Drug, and Cosmetic Act. <http://www.cfsan.fda.gov/~lrd/cfr110.html>
- 1034 Wachtel MR, Whitehand LC, and Mandrell RE. 2002a. Association of Escherichia coli O157:H7 with  
1035 preharvest leaf lettuce upon exposure to contaminated irrigation water. *Journal of food protection*  
1036 65 (1):18-25.  
1037 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11808792)  
1038 [uids=11808792](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11808792)
- 1039 Wachtel MR, Whitehand LC, and Mandrell RE. 2002b. Prevalence of Escherichia coli associated with a  
1040 cabbage crop inadvertently irrigated with partially treated sewage wastewater. *Journal of food protection*  
1041 65 (3):471-5.  
1042 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11899045)  
1043 [uids=11899045](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11899045)